



Introduction of Space Weather



Yihua Zheng

For SW REDI Winter 2014

- Space is NOT empty!
- There is weather in space - space weather
- Space is full of plasma - 4th state of matter (gas, liquid, solid)
 - plasma: important role in our plasma universe
 - plasma: important for space weather




Plasma




electrified gas with atoms dissociated into positive ions and negative electrons

- ☆ **99%** of the matter in universe is in the plasma state
 - ☆ Stellar interiors and atmospheres, gaseous nebulae, interstellar material are plasmas, Earth's ionosphere and above
- ☆ We live in the 1% of the universe in which plasmas do not occur naturally


A plasma is a quasineutral gas of charged and neutral particles which exhibits collective behavior



Prominence Eruption




Aurora: plasma interaction with molecules/atoms in the atmosphere



Plasma




A **plasma display** panel (PDP) is a type of flat panel display common to large TV displays 30 inches (76 cm) or larger. They are called "plasma" displays because the technology utilizes small cells containing electrically charged ionized gases, or what are in essence chambers more commonly known as fluorescent lamps.



Manmade: plasma TV, lamp, fluorescent tube, neon sign

Space Weather

- 
- ESA: “Space weather refers to the environmental conditions in Earth's magnetosphere, ionosphere and thermosphere due to the Sun and the solar wind that can influence the functioning and reliability of spaceborne and ground-based systems and services or endanger property or human health”.
- ESA: Space weather deals with phenomena involving ambient plasma, magnetic fields, radiation, particle flows in space and how these phenomena may influence man made systems. In addition to the Sun, non-solar sources such as galactic cosmic rays can be considered as space weather since they alter space environment conditions near the Earth.




Main Causes of Space Weather



- » **Solar Activity and the Solar Wind (Flare, CME, SEP, coronal hole high speed streams)**
- *driver*
- » **Galactic Cosmic Rays** - *driver*
- » **Meteoroids and Space Debris** - *driver*

The sun is the main driver of space weather



Main Causes of Space Weather



- » **Solar Activity and the Solar Wind** - *driver*
- » **Solar energetic particles and their entry into the magnetosphere** - *driver/internal response*
- » **Galactic Cosmic Rays** - *driver*
- » **Meteoroids and Space Debris** - *driver*
- » **The Earth's Magnetosphere: Geomagnetic Storms (and Substorms)** - *driver & internal response*
- » **Radiation Belts** - *driver & internal response*

Different regions of Space Weather






- Earth
- Mars
- Mercury
- Venus
- Spitzer
- Stereo_A
- Stereo_B

Orientation/Scales

1AU = 150 million km =
 93 million miles
 1 Rs = 695,500 km =
 110 RE
 1 RE = 6371 km
 MESSENGER: 0.3 AU
 1 Sun can fill a little
 more than 1 million
 Earth
 ACE/SOHO at L1: 1/100
 of the Sun-Earth
 distance



A movie on space weather


- http://missionscience.nasa.gov/sun/sunVideo_01spaceweather.html

- A movie on space weather

Space Weather Vocabulary


Why do we care?

Space Weather Illustrated



Sun and Earth are shown to approximate scale, but distance is not to scale.

The Sun maker of space weather




CME, Flares, and Coronal Hole HSS

Three very important solar wind disturbances/structures for space weather


- Radiation storm
- proton/ion radiation (SEP) <flare/CME>
- electron radiation <CIR HSS/CME>
- Radio blackout storm <flare>
- Geomagnetic storm
- CME storm (can be severe)
- CIR storm (moderate)

SWx impacts on spacecraft components


Major Space Environment Hazards




SWx impacts on satellite tracking




Satellite Tracking Problems After March 13-14, 1989 Storm





Main Drivers of Space weather: Flares/CMEs/ high speed solar wind streams



Solar Flares


radiation across the electromagnetic spectrum
most pronounced in EUV and soft X-ray

THE ELECTROMAGNETIC SPECTRUM


Penetrates Earth Atmosphere?




Wavelength (meters)




About the size of...




Frequency (Hz)



Temperature of bodies emitting the wavelength (K)



2012 March 7 X5.4/X1.3 Flares



Solar Flare:
sudden brightening on
the sun's surface




Flares radiate
throughout the
electromagnetic
spectrum

Most pronounced in x-
ray and EUV

Flare: SWx impacts

- Cause radio blackout through changing the structures/composition of the ionosphere (sudden ionospheric disturbances) – x ray and EUV emissions, lasting minutes to hours and dayside
- Affect radio comm., GPS, directly by its radio noises at different wavelengths
- Contribute to SEP – proton radiation, lasting a couple of days



Coronal Mass Ejections (CMEs)

CME


- ★ Massive burst of solar materials and magnetic field/flux into the interplanetary space: 10^{15} g
- ★ Kinetic energy 10^{32} erg
- ★ Yashiro et al. (2006) find that virtually all X-class flares have accompanying CMEs

CME viewed by coronagraph imagers




- ★ Eclipses allow corona to be better viewed
 - ★ Does not happen often
- ★ Modern coronagraph imager is inspired by that:
Occulting disk blocks the bright sun so we can observe corona features better

SEP: proton radiation




SWx Impacts of a CME

- ★ Contribute to SEP (proton/ion radiation): 20-30 minutes from the occurrence of the CME/flare
- ★ Result in a geomagnetic storm: takes 1-2 days arriving at Earth
- ★ Result in electron radiation enhancement in the near-Earth space (multiple CMEs): takes 1-3 days
- ★ Affecting spacecraft electronics – surfacing charging/internal charging, single event upsets (via SEPs)
- ★ Radio communication, navigation
- ★ Power grid, pipelines, and so on




Space Weather Effects and Timeline (Flare and CME)



Flare effects at Earth:
~ 8 minutes (radio blackout storms)
Duration: minutes to hours


SEP radiation effects reaching Earth: 20 minutes – 1 hour after the event onset
Duration: a few days

CME effects arrives @ Earth: 1-2 days (35 hours here)
Geomagnetic storms: a couple of days



Extras

Types of space weather events affecting nav and commu



UHF – GPS


- Energetic protons/ particles – via SEEs - affecting GPS satellites components
- Geomagnetic storms/ ionospheric storm - cause scintillations

VHF:



- Energetic protons - PCA
- Geomagnetic storms
- Solar radio emission associated with flare/CME

HF:


- Solar flares/x-ray
- Energetic protons - PCA
- Geomagnetic activities



CME and SEP path are different



CME: could get deflected, bended, but more or less in the radial direction



ENUL-2.7 lowres-2121-a3b1f WSA_V2.2 GONG-2121

ccmc/wsafr-oid/256x3

Important distinction

Ion Radiation storm vs Geomagnetic storm

CME impact and SEP (Solar Energetic Particle) impact are different

CME impact @ Earth:
Geomagnetic Storm

Radiation storm @ Earth from
SEPs

CME speed: 300 – 3500 km/s
SEPs: fraction of c
Light speed c: 3×10^5 km/s




SEPs: ion radiation storms

Potentially affect everywhere in the solar system




Courtesy: SVS@ NASA/GSFC

illustration of Geomagnetic storms due to a CME



Geomagnetic storms due to CIRs are at most moderate

- 
- The diagram shows the Earth with its magnetosphere, represented by blue, swirling magnetic field lines. The ionosphere and thermosphere are labeled in yellow text. The background is a dark space with stars.
1. SID (Sudden Ionospheric disturbance due to x-ray in solar flares)
dayside
 2. Solar energetic particle precipitation - particularly protons
High-latitude
 3. Geomagnetic storm disturbances
Ubiquitous/global

Eruptive solar events

Magnetosphere

**Communication/Navigation
Problem**

Solar radio bursts can directly affect GPS operation

- Solar radio bursts during December 2006 were sufficiently intense to be measurable with GPS receivers. The strongest event occurred on 6 December 2006 and affected the operation of many GPS receivers. This event exceeded 1,000,000 solar flux unit and was about 10 times larger than any previously reported event. The strength of the event was especially *surprising* since the solar radio bursts occurred near solar minimum. The strongest periods of solar radio burst activity lasted a few minutes to a few tens of minutes and, in some cases, exhibited large intensity differences between L1 (1575.42 MHz) and L2 (1227.60 MHz). Civilian dual frequency GPS receivers were the most severely affected, and these events suggest that continuous, precise positioning services should account for solar radio bursts in their operational plans. This investigation raises the possibility of even more intense solar radio bursts during the next solar maximum that will significantly impact the operation of GPS receivers.

Cerruti et al., 2008, Space Weather

Ionosphere Irregularities

- plasma bubbles: typical east–west dimensions of several hundred kilometers contain irregularities with scale-lengths ranging from tens of kilometers to tens of centimeters (Woodman and Tsunoda). Basu et al. (1978) showed that between sunset and midnight, 3-m scale irregularities that cause radar backscatter at 50 MHz, co-exist with sub-kilometer scale irregularities that cause VHF and L-band scintillations. After midnight, however, the radar backscatter and L-band scintillations decay but VHF scintillations caused by km-scale irregularities persist for several hours.